

Signatures of the Conformal Window

Elisabetta Pallante
University of Groningen

Work done in collaboration with:

A. Deuzeman, M.P. Lombardo, K. Miura, T. Nunes da Silva (lattice)

A. Barranco, J. Russo (AdS/CFT)

2008 $N_f=8$ is in the QCD phase

2009 $N_f=12$ is in the conformal window
(there is a conformal window)

Outline

Aims and strategy


The spectrum

The strong coupling side: Bulk transitions

The fate of $U(1)$ axial


Disappearance of conformality with AdS/CFT

Aims and Strategy I: Temperature - Flavor



Talk by K. Miura

Aims and Strategy II: Flavor - Coupling (T=0)



Miransky Yamawaki 1997

I Exact chiral $\beta < 0$	II Exact chiral $\beta > 0$	III Broken chiral UVFP?
----------------------------------	-----------------------------------	-------------------------------

The spectrum

QCD and non-QCD

$$\frac{m_\pi}{m_\rho} \sim \sqrt{m}$$

For fixed lattice spacing:


$$\frac{(am_\pi)^2}{am} \sim \text{const}$$

$$\frac{m_\pi}{m_\rho} \sim \text{const}$$

$$\frac{(am_\pi)^2}{am} \sim \frac{(am)^{2\delta \rightarrow 2}}{am} \sim (am)^{2\delta-1} \quad 0.5 < \delta \lesssim 1$$

Corrections to power laws are present for an interacting theory not at FP
(and finite volume)

The Edinburgh Plot of $N_f=12$ and $N_f=16$




Bare quark masses span a range 0.01 to 0.07 at various β for $N_f=12$

Bare quark masses span a range 0.025 to 0.15 at various β for $N_f=16$

Damgaard, Heller, Krasnitz, Olesen 1997

Fodor, Holland, Kuti, Nogradi, Schroeder 2011


QCD



↑ Ratio increases


This is compatible with a negative β function

and non-QCD




For a fixed m_π/m_ρ the inverted behavior with β_L is compatible with a positive β function

Masses and Power Laws




The ratio is approximately constant in a chirally symmetric phase. It goes to zero in the chiral (massless) limit of QCD.

The axial-vector mass splitting





Degeneracy in the chiral (massless) limit signals that chiral symmetry is restored.

Pseudo Goldstone mass and chiral condensate



Nf=12: lattice data




Exact chiral symmetry with non zero anomalous dimensions

Bulk transitions


Having already established that there is a bulk transition to a chirally broken phase...
Is it signalling the emergence of a UVFP? Or something else does happen.

The chiral condensate

Lattice fermion action: Naik improved staggered
Lattice gauge action: tree level Symanzik improved
Volumes: $16^3 \times 6$, $24^3 \times 8$, $24^3 \times 12$, $24^3 \times 24$ ($6^3 \times 6$, $12^3 \times 24$)
Lattice masses: 0.03, 0.025, 0.02, (0.015)



does not increase
with m decreasing



N_t dependence only for $N_t \leq 12$


The chiral cumulant

$$R = \chi_{\pi}^{-1} / \chi_{\sigma}^{-1}$$

$$\chi_{\sigma} = \chi_{\text{conn}} + \chi_{\text{disc}}$$

$$\chi_{\pi} = \langle \bar{\psi} \psi \rangle / m$$


$$M_p^2 = Z_p \chi_p^{-1}$$



$R \rightarrow 1$
 π - σ degeneracy

$R \rightarrow 0$
 π Goldstone

Zoom-in of chiral cumulant





$\leftrightarrow$



Inversion of mass ordering
at the chiral phase transition

The fate of $U(1)$ axial

Chiral susceptibilities mass dependence



Order parameter of U(1) axial



$SU(N_f)$ $\chi_\sigma - \chi_\pi = \chi_{\text{conn}} + \chi_{\text{disc}} - \chi_\pi \rightarrow \sigma$ - π degeneracy

$U_A(1)$ $\chi_\delta - \chi_\pi = \chi_{\text{conn}} - \chi_\pi \rightarrow \delta$ - π degeneracy

Summary: Signatures

The study of **symmetries and related phase transitions** in the relevant parameter space provides an ideal tool to explore theories inside and outside the CW.

Inside the CW, the **absence of (pseudo) Goldstone bosons** generates quite distinctive features as compared with QCD.


It is appealing the possibility that a **UVFP at strong coupling** emerges inside the CW, in addition to the IRFP. The first order nature of the chiral bulk transition excludes this possibility for $N_f=12$. However, a UVFP (second order PT) should appear at the end-point of a line of first order PTs --> end-point of the CW.

The appearance of an additional phase is plausibly related to the **partial restoration of $U(1)$ axial separately from the $SU(N_f)$ chiral restoration**. Additional effects such as an Aoki phase (staggered taste breaking) may add to this.

AdS/CFT


Disappearance of the CW

A new viable theory at Strong Coupling?



It suggests the disappearance of conformality
via annihilation of a pair of FPs

AdS/CFT



“IR/UV correspondence”

$z \rightarrow 0$	IR gravity
$z \rightarrow \infty$	UV field theory

Duality argument and exact β -function

Seiberg 1995

A conformal window for SQCD must be contained in the region
 $3/2 N_c < N_f < 3N_c$

NSVZ:
$$\beta_g = -\frac{g^3}{16\pi^2} \frac{3N_c - N_f(1 - \gamma_0)}{1 - \frac{g^2 N_c}{8\pi^2}}$$

FPS merging in “modified” SQCD?

Found a gravity solution that reproduces the NSVZ β -function of $\mathcal{N}=1$ SYM and confinement

Maldacena, Nunez 2004

Towards SQCD (large N_f, N_c : N_f/N_c fixed): SUGRA backgrounds

Massless case

Casero Nunez Paredes 2008

Massive case (smooth interpolation)

Conte Gaillard Ramallo 2011

$\Rightarrow$ SQCD + quartic operators

Mass interpolating function

$$0 \xleftarrow{0 \leftarrow r} N_f S(r) \xrightarrow{r \rightarrow \infty} N_f$$

Yang Mills
(Maldacena-Nunez solution)

Massless flavours
(linear dilaton)

The β -function from AdS/CFT

Ingredients: gravity solution - gauge coupling relation
energy - radius relation

Limitations: (gravity) singularities ($r \rightarrow 0$)

Recent results: Barranco EP Russo 2011

$N_f < 2N_c$ UV limit: $\beta \rightarrow \beta_{\text{NSVZ}}(\gamma_0 = -1/2)$
IR limit: ordinary confinement

$N_f = 2N_c$ **UVFP at strong coupling**

$N_f > 2N_c$ Seiberg dual ($N_c \rightarrow N_f - N_c$, $N_f - 2N_c$ flips sign)

Summary

Strongly coupled (quasi)conformal gauge theories might play a role in particle physics BSM

Large- N_f QCD is an instructive example

Conformal window opens around $N_f \sim 12$

Spectrum and phase transitions provide distinctive signatures of the conformal window

AdS/CFT suggests a unified view where 4D theories of particle physics are dual to gravity theories in AdS_5

AdS/CFT in phenomenology: warped RS1 models

AdS/CFT as a mathematical tool: FPs merging?

A UVFP is found at strong coupling for $N_f=2N_c$ in a class of gravity solutions dual to SQCD plus higher dimensional operators.