

Accelerating cosmology in modified gravity: what is the next?

Sergei D. Odintsov

Consejo Superior de Investigaciones Científicas, Institut de Ciències de l'Espai (ICE), (CSIC-IEEC), Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

KMI-IEEC Joint International Workshop, 3 August 2013

Contents

- 1 Introduction
- 2 $F(R)$ gravity
 - General properties
 - Scalar-tensor description
 - Viable modified gravities
- 3 $f(\mathcal{G})$ gravity
 - General properties
- 4 String-inspired model and scalar-Einstein-Gauss-Bonnet gravity
- 5 $F(R)$ bigravity
 - Cosmological Reconstruction and Cosmic Acceleration
- 6 What is the next?

Introduction

Accelerating cosmology within modified gravity: advances?

- 1 No need to introduce extra fields (inflaton, dark scalar or dark fluid, etc) to describe accelerating universe. The problem is solved by modification of gravitational action at early/late times!
- 2 Well-known applications to describe inflation in terms of higher-derivative gravity: Starobinsky, Mamaev-Mostepanenko, 1980.
- 3 Very natural possibility to describe dark energy era via modified gravity. The first discovery of quintessence dark era produced by power-law F(R) gravity is given by Capozziello (2002).
- 4 Very natural unification of inflation and dark energy eras in modified gravity: Nojiri-Odintsov 2003.
- 5 The complete description of the whole universe evolution eras sequence: inflation, radiation/matter dominance, dark energy in modified F(R) gravity, Nojiri-Odintsov 2006.
- 6 The possible emergence of dark matter from F(R) gravity (Capozziello 2004).
- 7 Direct relation of modified gravity theories with string theory. example of F(R) gravity (Nojiri-Odintsov 2003)
- 8 Relation with high energy physics (effective action, conformal anomaly, unification of GUTs with HD gravity)
- 9 Cosmological bounds and local tests.

Reviews:

Introduction to modified gravity and gravitational alternative for dark energy, Shin'ichi Nojiri (Japan, Natl. Defence Academy), Sergei D. Odintsov (ICREA, Barcelona and Barcelona, IECC). Jan 2006. 21 pp. Published in eConf C0602061 (2006) 06, Int.J.Geom.Meth.Mod.Phys. 4 (2007) 115-146, e-Print: hep-th/0601213

Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models, Shin'ichi Nojiri (Nagoya U. and KMI, Nagoya), Sergei D. Odintsov (ICREA, Barcelona and ICE, Bellaterra). Nov 2010. 98 pp. Published in Phys.Rept. 505 (2011) 59-148, e-Print: arXiv:1011.0544 [gr-qc]

Extended Theories of Gravity, Salvatore Capozziello, Mariafelicia De Laurentis (Naples U. and INFN, Naples). Aug 2011. 184 pp. Published in Phys.Rept. 509 (2011) 167-321, e-Print: arXiv:1108.6266

$F(R)$ gravity: General properties

The action of ghost-free $F(R)$ gravity

$$S_{F(R)} = \int d^4x \sqrt{-g} \left(\frac{F(R)}{2\kappa^2} + \mathcal{L}_{\text{matter}} \right). \quad (1)$$

The FRW equations in the Einstein gravity coupled with perfect fluid are given by

$$\rho_{\text{matter}} = \frac{3}{\kappa^2} H^2, \quad p_{\text{matter}} = -\frac{1}{\kappa^2} (3H^2 + 2\dot{H}), \quad (2)$$

which allow us to define an effective equation of state (EoS) parameter as follows:

$$w_{\text{eff}} = -1 - \frac{2\dot{H}}{3H^2}. \quad (3)$$

The field equation in the $F(R)$ gravity with matter is given by

$$\frac{1}{2} g_{\mu\nu} F(R) - R_{\mu\nu} F'(R) - g_{\mu\nu} \square F'(R) + \nabla_\mu \nabla_\nu F'(R) = -\frac{\kappa^2}{2} T_{\text{matter} \mu\nu}. \quad (4)$$

By assuming a spatially flat FRW universe,

$$ds^2 = -dt^2 + a(t)^2 \sum_{i=1,2,3} (dx^i)^2, \quad (5)$$

the equations corresponding to the FRW equations are given as follows:

$$0 = -\frac{F(R)}{2} + 3(H^2 + \dot{H}) F'(R) - 18(4H^2\dot{H} + H\ddot{H}) F''(R) + \kappa^2 \rho_{\text{matter}}, \quad (6)$$

$$0 = \frac{F(R)}{2} - (\dot{H} + 3H^2) F'(R) + 6(8H^2\dot{H} + 4\dot{H}^2 + 6H\ddot{H} + \ddot{H}) F''(R) \\ + 36(4H\dot{H} + \ddot{H})^2 F'''(R) + \kappa^2 p_{\text{matter}}. \quad (7)$$

F(R) gravity: General properties

One can find several (often exact) solutions of (6). When we neglect the contribution from matter, by assuming that the Ricci tensor is covariantly constant, that is, $R_{\mu\nu} \propto g_{\mu\nu}$, Eq. (4) reduces to an algebraic equation:

$$0 = 2F(R) - RF'(R). \quad (8)$$

If Eq. (8) has a solution, the (anti-)de Sitter, the Schwarzschild-(anti-)de Sitter space, and/or the Kerr-(anti-)de Sitter space is an exact vacuum solution.

Now we assume that $F(R)$ behaves as $F(R) \propto f_0 R^m$. Then Eq. (6) gives

$$0 = f_0 \left\{ -\frac{1}{2} (6\dot{H} + 12H^2)^m + 3m (\dot{H} + H^2) (6\dot{H} + 12H^2)^{m-1} - 3mH \frac{d}{dt} \left\{ (6\dot{H} + 12H^2)^{m-1} \right\} \right\} + \kappa^2 \rho_0 a^{-3(1+w)}. \quad (9)$$

Eq. (7) is irrelevant because it can be derived from (9). When the contribution from the matter can be neglected ($\rho_0 = 0$), the following solution exists:

$$H \sim \frac{-\frac{(m-1)(2m-1)}{m-2}}{t}, \quad (10)$$

which corresponds to the following EoS parameter (3):

$$w_{\text{eff}} = -\frac{6m^2 - 7m - 1}{3(m-1)(2m-1)}. \quad (11)$$

F(R) gravity: General properties

On the other hand, when the matter with a constant EoS parameter w is included, an exact solution of (9) is given by

$$a = a_0 t^{h_0}, \quad h_0 \equiv \frac{2m}{3(1+w)},$$

$$a_0 \equiv \left[-\frac{3f_0 h_0}{\kappa^2 \rho_0} \left(-6h_0 + 12h_0^2 \right)^{m-1} \{ (1-2m)(1-m) - (2-m)h_0 \} \right]^{-\frac{1}{3(1+w)}}, \quad (12)$$

and we find the effective EoS parameter (3) as

$$w_{\text{eff}} = -1 + \frac{w+1}{m}. \quad (13)$$

These solutions (10) and (12) show that modified gravity may describe early/late-time universe acceleration.

F(R) gravity: Scalar-tensor description

One can rewrite $F(R)$ gravity as the scalar-tensor theory. By introducing the auxiliary field A , the action (1) of the $F(R)$ gravity is rewritten in the following form:

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \{ F'(A) (R - A) + F(A) \} . \quad (14)$$

By the variation of A , one obtains $A = R$. Substituting $A = R$ into the action (14), one can reproduce the action in (1). Furthermore, by rescaling the metric as $g_{\mu\nu} \rightarrow e^\sigma g_{\mu\nu}$ ($\sigma = -\ln F'(A)$), we obtain the Einstein frame action:

$$S_E = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left(R - \frac{3}{2} g^{\rho\sigma} \partial_\rho \sigma \partial_\sigma \sigma - V(\sigma) \right) ,$$

$$V(\sigma) = e^\sigma g(e^{-\sigma}) - e^{2\sigma} f(g(e^{-\sigma})) = \frac{A}{F'(A)} - \frac{F(A)}{F'(A)^2} . \quad (15)$$

Here $g(e^{-\sigma})$ is given by solving the equation $\sigma = -\ln(1 + f'(A)) = -\ln F'(A)$ as $A = g(e^{-\sigma})$. Due to the conformal transformation, a coupling of the scalar field σ with usual matter arises. Since the mass of σ is given by

$$m_\sigma^2 \equiv \frac{3}{2} \frac{d^2 V(\sigma)}{d\sigma^2} = \frac{3}{2} \left\{ \frac{A}{F'(A)} - \frac{4F(A)}{(F'(A))^2} + \frac{1}{F''(A)} \right\} , \quad (16)$$

unless m_σ is very large, the large correction to the Newton law appears.

F(R) gravity: Viable modified gravities

As an example, we may consider the following exponential model

$$F(R) = R + \alpha \left(e^{-bR} - 1 \right). \quad (17)$$

Here α and b are constants. One can regard α as an effective cosmological constant and we choose the parameter b so that $1/b$ is much smaller than the curvature R_0 of the present universe. Then in the region $R \gg R_0$, we find

$$m_\sigma^2 \sim \frac{e^{bR}}{2\alpha b^2}, \quad (18)$$

which is positive and m_σ^2 could be very large and the correction to the Newton law is very small. In paper by Hu-Sawicki 2007, the one of the first examples of “realistic” $F(R)$ model was proposed. Currently, several viable models are proposed.

In order to obtain a realistic and viable model, $F(R)$ gravity should satisfy the following conditions:

- 1 When $R \rightarrow 0$, the Einstein gravity is recovered, that is,

$$F(R) \rightarrow R \quad \text{that is,} \quad \frac{F(R)}{R^2} \rightarrow \frac{1}{R}. \quad (19)$$

This also means that there is a flat space solution.

- 2 There appears a stable de Sitter solution, which corresponds to the late-time acceleration and, therefore, the curvature is small $R \sim R_L \sim (10^{-33} \text{ eV})^2$. This requires, when $R \sim R_L$,

$$\frac{F(R)}{R^2} = f_{0L} - f_{1L} (R - R_L)^{2n+2} + o\left((R - R_L)^{2n+2}\right). \quad (20)$$

Here, f_{0L} and f_{1L} are positive constants and n is a positive integer. Of course, in some cases this condition may not be strictly necessary.

F(R) gravity: Viable modified gravities

- 3 There appears a quasi-stable de Sitter solution that corresponds to the inflation of the early universe and, therefore, the curvature is large $R \sim R_I \sim (10^{16 \sim 19} \text{ GeV})^2$. The de Sitter space should not be exactly stable so that the curvature decreases very slowly. It requires

$$\frac{F(R)}{R^2} = f_{0I} - f_{1I} (R - R_I)^{2m+1} + o\left((R - R_I)^{2m+1}\right). \quad (21)$$

Here, f_{0I} and f_{1I} are positive constants and m is a positive integer.

- 4 In order to avoid the curvature singularity when $R \rightarrow \infty$, $F(R)$ should behaves as

$$F(R) \rightarrow f_\infty R^2 \quad \text{that is} \quad \frac{F(R)}{R^2} \rightarrow f_\infty. \quad (22)$$

Here, f_∞ is a positive and sufficiently small constant. Instead of (22), we may take

$$F(R) \rightarrow f_\infty R^{2-\epsilon} \quad \text{that is} \quad \frac{F(R)}{R^2} \rightarrow \frac{f_\infty}{R^\epsilon}. \quad (23)$$

Here, f_∞ is a positive constant and $0 < \epsilon < 1$. The above condition (22) or (23) prevents both the future singularity and the singularity due to large density of matter.

- 5 To avoid the anti-gravity, we require

$$F'(R) > 0, \quad (24)$$

which is rewritten as

$$\frac{d}{dR} \left(\ln \left(\frac{F(R)}{R^2} \right) \right) - \frac{2}{R}. \quad (25)$$

F(R) gravity: Viable modified gravities

- ⑥ Combining conditions (19) and (24), one finds

$$F(R) > 0. \quad (26)$$

- ⑦ To avoid the matter instability (Dolgov-Kawasaki 2003), we require

$$U(R_b) \equiv \frac{R_b}{3} - \frac{F^{(1)}(R_b)F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2} - \frac{F^{(1)}(R_b)}{3F^{(2)}(R_b)} \\ + \frac{2F(R_b)F^{(3)}(R_b)}{3F^{(2)}(R_b)^2} - \frac{F^{(3)}(R_b)R_b}{3F^{(2)}(R_b)^2} < 0. \quad (27)$$

The conditions 1 and 2 tell that an extra, unstable de Sitter solution must appear at $R = R_e$ ($0 < R_e < R_L$). Since the universe evolution will stop at $R = R_L$ because the de Sitter solution $R = R_L$ is stable; the curvature never becomes smaller than R_L and, therefore, the extra de Sitter solution is not realized.

An example of viable $F(R)$ gravity is given below

$$\frac{F(R)}{R^2} = \left\{ (X_m(R_i; R) - X_m(R_i; R_1))(X_m(R_i; R) - X_m(R_i; R_L))^{2n+2} \right. \\ \left. + X_m(R_i; R_1)X_m(R_i; R_L)^{2n+2} + f_\infty^{2n+3} \right\}^{\frac{1}{2n+3}}, \\ X_m(R_i; R) \equiv \frac{(2m+1)R_i^{2m}}{(R-R_i)^{2m+1} + R_i^{2m+1}}. \quad (28)$$

Here, n and m are integers greater or equal to unity, and $n, m \geq 1$ and R_1 is a parameter related with R_e by

$$X(R_i; R_e) = \frac{(2n+2)X(R_i; R_1)X(R_i; R_1) + X(R_i; R_L)}{2n+3}. \quad (29)$$

We also assume $0 < R_1 < R_L \ll R_i$.

F(R) gravity: Viable modified gravities

Another realistic theory unifying inflation with dark energy is given in

$$F(R) = R - 2\Lambda \left(1 - e^{-\frac{R}{R_0}}\right) - \Lambda_i \left(1 - e^{-\left(\frac{R}{R_i}\right)^n}\right) + \gamma R^\alpha . \quad (30)$$

Here Λ is the effective cosmological constant in the present universe and we also assume the parameter R_0 is almost equal to Λ . R_i and Λ_i are typical values of the curvature and the effective cosmological constant. α is a constant: $1 < \alpha \leq 2$. Generalizations: coupling of curvature with trace of EMT (Harko-Lobo- -Nojiri-Odintsov) or with EMT (Saez-Gomez).

f(G) gravity: General properties

Topological Gauss-Bonnet invariant:

$$\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\xi\sigma}R^{\mu\nu\xi\sigma}, \quad (31)$$

is added to the action of the Einstein gravity. One starts with the following action:

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R + f(\mathcal{G}) + \mathcal{L}_{\text{matter}} \right). \quad (32)$$

Here, $\mathcal{L}_{\text{matter}}$ is the Lagrangian density of matter. The variation of the metric $g_{\mu\nu}$:

$$\begin{aligned} 0 = & \frac{1}{2\kappa^2} \left(-R^{\mu\nu} + \frac{1}{2}g^{\mu\nu}R \right) + T_{\text{matter}}^{\mu\nu} + \frac{1}{2}g^{\mu\nu}f(\mathcal{G}) - 2f'(\mathcal{G})RR^{\mu\nu} \\ & + 4f'(\mathcal{G})R^\mu_\rho R^{\nu\rho} - 2f'(\mathcal{G})R^{\mu\rho\sigma\tau}R^\nu_{\rho\sigma\tau} - 4f'(\mathcal{G})R^{\mu\rho\sigma\nu}R_{\rho\sigma} + 2(\nabla^\mu\nabla^\nu f'(\mathcal{G}))R \\ & - 2g^{\mu\nu}(\nabla^2 f'(\mathcal{G}))R - 4(\nabla_\rho\nabla^\mu f'(\mathcal{G}))R^{\nu\rho} - 4(\nabla_\rho\nabla^\nu f'(\mathcal{G}))R^{\mu\rho} \\ & + 4(\nabla^2 f'(\mathcal{G}))R^{\mu\nu} + 4g^{\mu\nu}(\nabla_\rho\nabla_\sigma f'(\mathcal{G}))R^{\rho\sigma} - 4(\nabla_\rho\nabla_\sigma f'(\mathcal{G}))R^{\mu\rho\nu\sigma}. \end{aligned} \quad (33)$$

The first FRW equation:

$$0 = -\frac{3}{\kappa^2}H^2 - f(\mathcal{G}) + \mathcal{G}f'(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3 + \rho_{\text{matter}}. \quad (34)$$

Here \mathcal{G} has the following form:

$$\mathcal{G} = 24 \left(H^2 \dot{H} + H^4 \right). \quad (35)$$

the FRW-like equations (fluid description):

$$\rho_{\text{eff}}^{\mathcal{G}} = \frac{3}{\kappa^2}H^2, \quad p_{\text{eff}}^{\mathcal{G}} = -\frac{1}{\kappa^2} \left(3H^2 + 2\dot{H} \right). \quad (36)$$

f(G) gravity: General properties

Here,

$$\rho_{\text{eff}}^{\mathcal{G}} \equiv -f(\mathcal{G}) + \mathcal{G}f'(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3 + \rho_{\text{matter}},$$

$$p_{\text{eff}}^{\mathcal{G}} \equiv f(\mathcal{G}) - \mathcal{G}f'(\mathcal{G}) + \frac{2\mathcal{G}\dot{\mathcal{G}}}{3H}f''(\mathcal{G}) + 8H^2\ddot{\mathcal{G}}f''(\mathcal{G}) + 8H^2\dot{\mathcal{G}}^2f'''(\mathcal{G}) + p_{\text{matter}}. \quad (37)$$

When $\rho_{\text{matter}} = 0$, Eq. (34) has a de Sitter universe solution where H , and therefore \mathcal{G} , are constant. For $H = H_0$, with a constant H_0 , Eq. (34) turns into

$$0 = -\frac{3}{\kappa^2}H_0^2 + 24H_0^4f'(24H_0^4) - f(24H_0^4). \quad (38)$$

As an example, we consider the model

$$f(\mathcal{G}) = f_0 |\mathcal{G}|^\beta, \quad (39)$$

with constants f_0 and β . Then, the solution of Eq. (38) is given by

$$H_0^4 = \frac{1}{24(8(n-1)\kappa^2f_0)^{\frac{1}{\beta-1}}}. \quad (40)$$

No matter and GR. Eq. (34) reduces to

$$0 = \mathcal{G}f'(\mathcal{G}) - f(\mathcal{G}) - 24\dot{\mathcal{G}}f''(\mathcal{G})H^3. \quad (41)$$

If $f(\mathcal{G})$ behaves as (39), assuming

$$a = \begin{cases} a_0 t^{h_0} & \text{when } h_0 > 0 \text{ (quintessence)} \\ a_0 (t_s - t)^{h_0} & \text{when } h_0 < 0 \text{ (phantom)} \end{cases}, \quad (42)$$

one obtains

$$0 = (\beta - 1)h_0^6(h_0 - 1)(h_0 - 1 + 4\beta). \quad (43)$$

f(G) gravity: General properties

As $h_0 = 1$ implies $\mathcal{G} = 0$, one may choose

$$h_0 = 1 - 4\beta, \quad (44)$$

and Eq. (3) gives

$$w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta)}. \quad (45)$$

Therefore, if $\beta > 0$, the universe is accelerating ($w_{\text{eff}} < -1/3$), and if $\beta > 1/4$, the universe is in a phantom phase ($w_{\text{eff}} < -1$). Thus, we are led to consider the following model:

$$f(\mathcal{G}) = f_i |\mathcal{G}|^{\beta_i} + f_l |\mathcal{G}|^{\beta_l}, \quad (46)$$

where it is assumed that

$$\beta_i > \frac{1}{2}, \quad \frac{1}{2} > \beta_l > \frac{1}{4}. \quad (47)$$

Then, when the curvature is large, as in the primordial universe, the first term dominates, compared with the second term and the Einstein term, and it gives

$$-1 > w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta_i)} > -\frac{5}{3}. \quad (48)$$

On the other hand, when the curvature is small, as is the case in the present universe, the second term in (46) dominates compared with the first term and the Einstein term and yields

$$w_{\text{eff}} = -1 + \frac{2}{3(1 - 4\beta_l)} < -\frac{5}{3}. \quad (49)$$

Therefore, theory (46) can produce a model that is able to describe inflation and the late-time acceleration of the universe in a unified manner.

f(G) gravity: General properties

The action (32) can be rewritten by introducing the auxiliary scalar field ϕ as,

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - V(\phi) - \xi(\phi)\mathcal{G} \right]. \quad (50)$$

By variation over ϕ , one obtains

$$0 = V'(\phi) + \xi'(\phi)\mathcal{G}, \quad (51)$$

which could be solved with respect to ϕ as

$$\phi = \phi(\mathcal{G}). \quad (52)$$

By substituting the expression (52) into the action (50), we obtain the action of f(G) gravity, with

$$f(\mathcal{G}) = -V(\phi(\mathcal{G})) + \xi(\phi(\mathcal{G}))\mathcal{G}. \quad (53)$$

Assuming a spatially-flat FRW universe and the scalar field ϕ to depend only on t , we obtain the field equations:

$$0 = -\frac{3}{\kappa^2}H^2 + V(\phi) + 24H^3 \frac{d\xi(\phi(t))}{dt}, \quad (54)$$

$$0 = \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) - V(\phi) - 8H^2 \frac{d^2\xi(\phi(t))}{dt^2} - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} - 16H^3 \frac{d\xi(\phi(t))}{dt}. \quad (55)$$

Combining the above equations, we obtain

$$\begin{aligned} 0 &= \frac{2}{\kappa^2} \dot{H} - 8H^2 \frac{d^2\xi(\phi(t))}{dt^2} - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} + 8H^3 \frac{d\xi(\phi(t))}{dt} \\ &= \frac{2}{\kappa^2} \dot{H} - 8a \frac{d}{dt} \left(\frac{H^2}{a} \frac{d\xi(\phi(t))}{dt} \right), \end{aligned} \quad (56)$$

f(G) gravity: General properties

which can be solved with respect to $\xi(\phi(t))$ as

$$\xi(\phi(t)) = \frac{1}{8} \int^t dt_1 \frac{a(t_1)}{H(t_1)^2} W(t_1), \quad W(t) \equiv \frac{2}{\kappa^2} \int^t \frac{dt_1}{a(t_1)} \dot{H}(t_1). \quad (57)$$

Combining (54) and (57), the expression for $V(\phi(t))$ follows:

$$V(\phi(t)) = \frac{3}{\kappa^2} H(t)^2 - 3a(t)H(t)W(t). \quad (58)$$

As there is a freedom of redefinition of the scalar field ϕ , we may identify t with ϕ . Hence, we consider the model where $V(\phi)$ and $\xi(\phi)$ can be expressed in terms of a single function g as

$$\begin{aligned} V(\phi) &= \frac{3}{\kappa^2} g'(\phi)^2 - 3g'(\phi) e^{g(\phi)} U(\phi), \\ \xi(\phi) &= \frac{1}{8} \int^\phi d\phi_1 \frac{e^{g(\phi_1)}}{g'(\phi_1)^2} U(\phi_1), \\ U(\phi) &\equiv \frac{2}{\kappa^2} \int^\phi d\phi_1 e^{-g(\phi_1)} g''(\phi_1). \end{aligned} \quad (59)$$

By choosing $V(\phi)$ and $\xi(\phi)$ as (59), one can easily find the following solution for Eqs.(54) and (55):

$$a = a_0 e^{g(t)} \quad (H = g'(t)). \quad (60)$$

Therefore one can reconstruct $F(G)$ gravity to generate arbitrary expansion history of the universe. Thus, we reviewed the modified Gauss-Bonnet gravity and demonstrated that it may naturally lead to the unified cosmic history, including the inflation and dark energy era.

String-inspired model and scalar-Einstein-Gauss-Bonnet gravity

Stringy gravity:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2} + \mathcal{L}_\phi + \mathcal{L}_c + \dots \right], \quad (61)$$

where ϕ is the dilaton, \mathcal{L}_ϕ is the Lagrangian of ϕ , and \mathcal{L}_c expresses the string curvature correction terms,

$$\mathcal{L}_\phi = -\partial_\mu \phi \partial^\mu \phi - V(\phi), \quad \mathcal{L}_c = c_1 \alpha' e^{2\frac{\phi}{\phi_0}} \mathcal{L}_c^{(1)} + c_2 \alpha'^2 e^{4\frac{\phi}{\phi_0}} \mathcal{L}_c^{(2)} + c_3 \alpha'^3 e^{6\frac{\phi}{\phi_0}} \mathcal{L}_c^{(3)}, \quad (62)$$

where $1/\alpha'$ is the string tension, $\mathcal{L}_c^{(1)}$, $\mathcal{L}_c^{(2)}$, and $\mathcal{L}_c^{(3)}$ express the leading-order (Gauss-Bonnet term \mathcal{G} in (31)), the second-order, and the third-order curvature corrections, respectively:

$$\mathcal{L}_c^{(1)} = \Omega_2, \quad \mathcal{L}_c^{(2)} = 2\Omega_3 + R_{\alpha\beta}^{\mu\nu} R_{\lambda\rho}^{\alpha\beta} R_{\mu\nu}^{\lambda\rho}, \quad \mathcal{L}_c^{(3)} = \mathcal{L}_{31} - \delta_H \mathcal{L}_{32} - \frac{\delta_B}{2} \mathcal{L}_{33}. \quad (63)$$

Here, δ_B and δ_H take the value of 0 or 1 and

$$\Omega_2 = \mathcal{G},$$

$$\Omega_3 \propto \epsilon^{\mu\nu\rho\sigma\tau\eta} \epsilon_{\mu'\nu'\rho'\sigma'\tau'\eta'} R_{\mu\nu}^{\mu'\nu'} R_{\rho\sigma}^{\rho'\sigma'} R_{\tau\eta}^{\tau'\eta'},$$

$$\mathcal{L}_{31} = \zeta(3) R_{\mu\nu\rho\sigma} R^{\alpha\nu\rho\beta} \left(R_{\delta\beta}^{\mu\gamma} R_{\alpha\gamma}^{\delta\sigma} - 2R_{\delta\alpha}^{\mu\gamma} R_{\beta\gamma}^{\delta\sigma} \right),$$

$$\mathcal{L}_{32} = \frac{1}{8} \left(R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} \right)^2 + \frac{1}{4} R_{\mu\nu}^{\gamma\delta} R_{\gamma\delta}^{\rho\sigma} R_{\rho\sigma}^{\alpha\beta} R_{\alpha\beta}^{\mu\nu} - \frac{1}{2} R_{\mu\nu}^{\alpha\beta} R_{\alpha\beta}^{\rho\sigma} R_{\sigma\gamma\delta}^{\mu} R_{\rho}^{\nu\gamma\delta} - R_{\mu\nu}^{\alpha\beta} R_{\alpha\beta}^{\rho\nu} R_{\rho\sigma}^{\gamma\delta} R_{\gamma}^{\sigma}$$

$$\mathcal{L}_{33} = \left(R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} \right)^2 - 10 R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\sigma} R_{\sigma\gamma\delta\rho} R^{\beta\gamma\delta\rho} - R_{\mu\nu\alpha\beta} R^{\mu\nu\rho\sigma} R^{\beta\sigma\gamma\delta} R_{\delta\gamma\rho}^{\alpha} \quad (64)$$

The correction terms are different depending on the type of string theory; the dependence is encoded in the curvature invariants and in the coefficients (c_1, c_2, c_3) and δ_H, δ_B , as follows,

- For the Type II superstring theory: $(c_1, c_2, c_3) = (0, 0, 1/8)$ and $\delta_H = \delta_B = 0$.
- For the heterotic superstring theory: $(c_1, c_2, c_3) = (1/8, 0, 1/8)$ and $\delta_H = 1, \delta_B = 0$.
- For the bosonic superstring theory: $(c_1, c_2, c_3) = (1/4, 1/48, 1/8)$ and $\delta_H = 0, \delta_B = 1$.

String-inspired model and scalar-Einstein-Gauss-Bonnet gravity

The starting action is:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa^2} - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - V(\phi) - \xi(\phi) \mathcal{G} \right]. \quad (65)$$

Field equations:

$$\begin{aligned} 0 = & \frac{1}{\kappa^2} \left(-R^{\mu\nu} + \frac{1}{2} g^{\mu\nu} R \right) + \frac{1}{2} \partial^\mu \phi \partial^\nu \phi - \frac{1}{4} g^{\mu\nu} \partial_\rho \phi \partial^\rho \phi + \frac{1}{2} g^{\mu\nu} (-V(\phi) + \xi(\phi) \mathcal{G}) \\ & - 2\xi(\phi) R R^{\mu\nu} - 4\xi(\phi) R^\mu{}_\rho R^{\nu\rho} - 2\xi(\phi) R^{\mu\rho\sigma\tau} R^\nu{}_{\rho\sigma\tau} + 4\xi(\phi) R^{\mu\rho\nu\sigma} R_{\rho\sigma} \\ & + 2 (\nabla^\mu \nabla^\nu \xi(\phi)) R - 2g^{\mu\nu} (\nabla^2 \xi(\phi)) R - 4 (\nabla_\rho \nabla^\mu \xi(\phi)) R^{\nu\rho} - 4 (\nabla_\rho \nabla^\nu \xi(\phi)) R^{\mu\rho} \\ & + 4 (\nabla^2 \xi(\phi)) R^{\mu\nu} + 4g^{\mu\nu} (\nabla_\rho \nabla_\sigma \xi(\phi)) R^{\rho\sigma} + 4 (\nabla_\rho \nabla_\sigma \xi(\phi)) R^{\mu\rho\nu\sigma}. \end{aligned} \quad (66)$$

FRW eq.:

$$0 = -\frac{3}{\kappa^2} H^2 + \frac{1}{2} \dot{\phi}^2 + V(\phi) + 24H^3 \frac{d\xi(\phi(t))}{dt}, \quad (67)$$

$$\begin{aligned} 0 = & \frac{1}{\kappa^2} \left(2\dot{H} + 3H^2 \right) + \frac{1}{2} \dot{\phi}^2 - V(\phi) - 8H^2 \frac{d^2 \xi(\phi(t))}{dt^2} \\ & - 16H\dot{H} \frac{d\xi(\phi(t))}{dt} - 16H^3 \frac{d\xi(\phi(t))}{dt}. \end{aligned} \quad (68)$$

Scalar equation

$$0 = \ddot{\phi} + 3H\dot{\phi} + V'(\phi) + \xi'(\phi) \mathcal{G}. \quad (69)$$

String-inspired model and scalar-Einstein-Gauss-Bonnet gravity

In particular when we consider the following string-inspired model,

$$V = V_0 e^{-\frac{2\phi}{\phi_0}}, \quad \xi(\phi) = \xi_0 e^{\frac{2\phi}{\phi_0}}, \quad (70)$$

the de Sitter space solution follows:

$$H^2 = H_0^2 \equiv -\frac{e^{-\frac{2\phi_0}{\phi_0}}}{8\xi_0\kappa^2}, \quad \phi = \phi_0. \quad (71)$$

Here, ϕ_0 is an arbitrary constant. If ϕ_0 is chosen to be larger, the Hubble rate $H = H_0$ becomes smaller. Then, if $\xi_0 \sim \mathcal{O}(1)$, by choosing $\phi_0/\phi_0 \sim 140$, the value of the Hubble rate $H = H_0$ is consistent with the observations. The model (70) also has another solution:

$$\begin{aligned} H &= \frac{h_0}{t}, \quad \phi = \phi_0 \ln \frac{t}{t_1} && \text{when } h_0 > 0, \\ H &= -\frac{h_0}{t_s - t}, \quad \phi = \phi_0 \ln \frac{t_s - t}{t_1} && \text{when } h_0 < 0. \end{aligned} \quad (72)$$

Here, h_0 is obtained by solving the following algebraic equations:

$$0 = -\frac{3h_0^2}{\kappa^2} + \frac{\phi_0^2}{2} + V_0 t_1^2 - \frac{48\xi_0 h_0^3}{t_1^2}, \quad 0 = (1 - 3h_0)\phi_0^2 + 2V_0 t_1^2 + \frac{48\xi_0 h_0^3}{t_1^2} (h_0 - 1). \quad (73)$$

Eqs. (73) can be rewritten as

$$\begin{aligned} V_0 t_1^2 &= -\frac{1}{\kappa^2(1+h_0)} \left\{ 3h_0^2(1-h_0) + \frac{\phi_0^2 \kappa^2 (1-5h_0)}{2} \right\}, && (74) \\ \frac{48\xi_0 h_0^2}{t_1^2} &= -\frac{6}{\kappa^2(1+h_0)} \left(h_0 - \frac{\phi_0^2 \kappa^2}{2} \right). && (75) \end{aligned}$$

The arbitrary value of h_0 can be realized by properly choosing V_0 and ξ_0 . With the appropriate choice of V_0 and ξ_0 , we can obtain a negative h_0 and, therefore, the effective EoS parameter (3) is less than -1 , $w_{\text{eff}} < -1$, which corresponds to the effective phantom.

F(R) bigravity

Non-linear massive gravity (with non-dynamical background metric) was extended to the ghost-free construction with the dynamical metric (Hassan et al).

The convenient description of the theory gives bigravity or bimetric gravity which contains two metrics (symmetric tensor fields). One of two metrics is called physical metric while second metric is called reference metric.

Next is $F(R)$ bigravity which is also ghost-free theory. We introduce four kinds of metrics, $g_{\mu\nu}$, $g_{\mu\nu}^J$, $f_{\mu\nu}$, and $f_{\mu\nu}^J$. The physical observable metric $g_{\mu\nu}^J$ is the metric in the Jordan frame. The metric $g_{\mu\nu}$ corresponds to the metric in the Einstein frame in the standard $F(R)$ gravity and therefore the metric $g_{\mu\nu}$ is not physical metric. In the bigravity theories, we have to introduce another reference metrics or symmetric tensor $f_{\mu\nu}$ and $f_{\mu\nu}^J$. The metric $f_{\mu\nu}$ is the metric corresponding to the Einstein frame with respect to the curvature given by the metric $f_{\mu\nu}$. On the other hand, the metric $f_{\mu\nu}^J$ is the metric corresponding to the Jordan frame.

The starting action is given by

$$S_{\text{bi}} = M_g^2 \int d^4x \sqrt{-\det g} R^{(g)} + M_f^2 \int d^4x \sqrt{-\det f} R^{(f)} + 2m^2 M_{\text{eff}}^2 \int d^4x \sqrt{-\det g} \sum_{n=0}^4 \beta_n e_n \left(\sqrt{g^{-1}f} \right). \quad (77)$$

Here $R^{(g)}$ is the scalar curvature for $g_{\mu\nu}$ and $R^{(f)}$ is the scalar curvature for $f_{\mu\nu}$. M_{eff} is defined by

$$\frac{1}{M_{\text{eff}}^2} = \frac{1}{M_g^2} + \frac{1}{M_f^2}. \quad (78)$$

Furthermore, tensor $\sqrt{g^{-1}f}$ is defined by the square root of $g^{\mu\rho} f_{\rho\nu}$, that is, $\left(\sqrt{g^{-1}f} \right)_{\rho}^{\mu} \left(\sqrt{g^{-1}f} \right)_{\nu}^{\rho} = g^{\mu\rho} f_{\rho\nu}$.

F(R) bigravity

For general tensor X^μ_ν , $e_n(X)$'s are defined by

$$\begin{aligned} e_0(X) &= 1, & e_1(X) &= [X], & e_2(X) &= \frac{1}{2}([X]^2 - [X^2]), \\ e_3(X) &= \frac{1}{6}([X]^3 - 3[X][X^2] + 2[X^3]), \\ e_4(X) &= \frac{1}{24}([X]^4 - 6[X]^2[X^2] + 3[X^2]^2 + 8[X][X^3] - 6[X^4]), \\ e_k(X) &= 0 \text{ for } k > 4. \end{aligned} \quad (79)$$

Here $[X]$ expresses the trace of arbitrary tensor X^μ_ν : $[X] = X^\mu_\mu$. In order to construct the consistent F(R) bigravity, we add the following terms to the action (77):

$$S_\varphi = -M_g^2 \int d^4x \sqrt{-\det g} \left\{ \frac{3}{2} g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi + V(\varphi) \right\} + \int d^4x \mathcal{L}_{\text{matter}}(e^\varphi g_{\mu\nu}, \Phi_i), \quad (80)$$

$$S_\xi = -M_f^2 \int d^4x \sqrt{-\det f} \left\{ \frac{3}{2} f^{\mu\nu} \partial_\mu \xi \partial_\nu \xi + U(\xi) \right\}. \quad (81)$$

By the conformal transformations $g_{\mu\nu} \rightarrow e^{-\varphi} g_{\mu\nu}^J$ and $f_{\mu\nu} \rightarrow e^{-\xi} f_{\mu\nu}^J$, the total action $S_F = S_{\text{bi}} + S_\varphi + S_\xi$ is transformed as

$$\begin{aligned} S_F &= M_f^2 \int d^4x \sqrt{-\det f^J} \left\{ e^{-\xi} R^{J(f)} - e^{-2\xi} U(\xi) \right\} \\ &+ 2m^2 M_{\text{eff}}^2 \int d^4x \sqrt{-\det g^J} \sum_{n=0}^4 \beta_n e^{(\frac{n}{2}-2)\varphi - \frac{n}{2}\xi} e_n \left(\sqrt{g^{J-1} f^J} \right) \\ &+ M_g^2 \int d^4x \sqrt{-\det g^J} \left\{ e^{-\varphi} R^{J(g)} - e^{-2\varphi} V(\varphi) \right\} \\ &+ \int d^4x \mathcal{L}_{\text{matter}}(g_{\mu\nu}^J, \Phi_i). \end{aligned} \quad (82)$$

F(R) bigravity

The kinetic terms for φ and ξ vanish. By the variations with respect to φ and ξ as in the case of convenient $F(R)$ gravity, we obtain

$$0 = 2m^2 M_{\text{eff}}^2 \sum_{n=0}^4 \beta_n \left(\frac{n}{2} - 2 \right) e^{\left(\frac{n}{2}-2\right)\varphi - \frac{n}{2}\xi} e_n \left(\sqrt{g^{J-1} f^J} \right) + M_g^2 \left\{ -e^{-\varphi} R^{J(g)} + 2e^{-2\varphi} V(\varphi) + e^{-2\varphi} V'(\varphi) \right\}, \quad (83)$$

$$0 = -2m^2 M_{\text{eff}}^2 \sum_{n=0}^4 \frac{\beta_n n}{2} e^{\left(\frac{n}{2}-2\right)\varphi - \frac{n}{2}\xi} e_n \left(\sqrt{g^{J-1} f^J} \right) + M_f^2 \left\{ -e^{-\xi} R^{J(f)} + 2e^{-2\xi} U(\xi) + e^{-2\xi} U'(\xi) \right\}. \quad (84)$$

The Eqs. (83) and (84) can be solved algebraically with respect to φ and ξ as

$$\varphi = \varphi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)$$

and

$$\xi = \xi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)$$

. Substituting above φ and ξ into (82), one gets $F(R)$ bigravity:

$$S_F = M_f^2 \int d^4x \sqrt{-\det f^J} F^{(f)} \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) + 2m^2 M_{\text{eff}}^2 \int d^4x \sqrt{-\det g} \sum_{n=0}^4 \beta_n e^{\left(\frac{n}{2}-2\right)\varphi \left(R^{J(g)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)} e_n \left(\sqrt{g^{J-1} f^J} \right) + M_g^2 \int d^4x \sqrt{-\det g^J} F^{J(g)} \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) + \int d^4x \mathcal{L}_{\text{matter}} \left(g_{\mu\nu}^J, \Phi_i \right), \quad (85)$$

F(R) bigravity

$$F^{J(g)} \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) \equiv \left\{ e^{-\varphi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)} R^{J(g)} \right. \\ \left. - e^{-2\varphi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)} V \left(\varphi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) \right) \right\}, \quad (86)$$

$$F^{(f)} \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) \equiv \left\{ e^{-\xi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)} R^{J(f)} \right. \\ \left. - e^{-2\xi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right)} U \left(\xi \left(R^{J(g)}, R^{J(f)}, e_n \left(\sqrt{g^{J-1} f^J} \right) \right) \right) \right\}. \quad (87)$$

Note that it is difficult to solve Eqs. (83) and (84) with respect to φ and ξ explicitly. Therefore, it might be easier to define the model in terms of the auxiliary scalars φ and ξ as in (82).

F(R) bigravity: Cosmological Reconstruction and Cosmic Acceleration

Let us consider the cosmological reconstruction program. For simplicity, we start from the minimal case

$$S_{\text{bi}} = M_g^2 \int d^4x \sqrt{-\det g} R^{(g)} + M_f^2 \int d^4x \sqrt{-\det f} R^{(f)} + 2m^2 M_{\text{eff}}^2 \int d^4x \sqrt{-\det g} \left(3 - \text{tr} \sqrt{g^{-1}f} + \det \sqrt{g^{-1}f} \right). \quad (88)$$

In order to evaluate $\delta \sqrt{g^{-1}f}$, two matrices M and N , which satisfy the relation $M^2 = N$ are taken. Since $\delta MM + M\delta M = \delta N$, one finds

$$\text{tr} \delta M = \frac{1}{2} \text{tr} \left(M^{-1} \delta N \right). \quad (89)$$

For a while, we consider the Einstein frame action (88) with (80) and (81) but matter contribution is neglected. Then by the variation over $g_{\mu\nu}$, we obtain

$$0 = M_g^2 \left(\frac{1}{2} g_{\mu\nu} R^{(g)} - R_{\mu\nu}^{(g)} \right) + m^2 M_{\text{eff}}^2 \left\{ g_{\mu\nu} \left(3 - \text{tr} \sqrt{g^{-1}f} \right) + \frac{1}{2} f_{\mu\rho} \left(\sqrt{g^{-1}f} \right)^{-1\rho}{}_{\nu} + \frac{1}{2} f_{\nu\rho} \left(\sqrt{g^{-1}f} \right)^{-1\rho}{}_{\mu} \right\} + M_g^2 \left[\frac{1}{2} \left(\frac{3}{2} g^{\rho\sigma} \partial_\rho \varphi \partial_\sigma \varphi + V(\varphi) \right) g_{\mu\nu} - \frac{3}{2} \partial_\mu \varphi \partial_\nu \varphi \right]. \quad (90)$$

On the other hand, by the variation over $f_{\mu\nu}$, we get

$$0 = M_f^2 \left(\frac{1}{2} f_{\mu\nu} R^{(f)} - R_{\mu\nu}^{(f)} \right) + m^2 M_{\text{eff}}^2 \sqrt{\det(f^{-1}g)} \left\{ -\frac{1}{2} f_{\mu\rho} \left(\sqrt{g^{-1}f} \right)^\rho{}_{\nu} - \frac{1}{2} f_{\nu\rho} \left(\sqrt{g^{-1}f} \right)^\rho{}_{\mu} + \det \left(\sqrt{g^{-1}f} \right) f_{\mu\nu} \right\} + M_f^2 \left[\frac{1}{2} \left(\frac{3}{2} f^{\rho\sigma} \partial_\rho \xi \partial_\sigma \xi + U(\xi) \right) f_{\mu\nu} - \frac{3}{2} \partial_\mu \xi \partial_\nu \xi \right]. \quad (91)$$

F(R) bigravity: Cosmological Reconstruction and Cosmic Acceleration

We should note that $\det \sqrt{g} \det \sqrt{g^{-1}f} \neq \sqrt{\det f}$ in general. The variations of the scalar fields φ and ξ are given by

$$0 = -3\Box_g \varphi + V'(\varphi), \quad 0 = -3\Box_f \xi + U'(\xi). \quad (92)$$

Here \Box_g (\Box_f) is the d'Alembertian with respect to the metric g (f). By multiplying the covariant derivative ∇_g^μ with respect to the metric g with Eq. (90) and using the Bianchi identity $0 = \nabla_g^\mu \left(\frac{1}{2} g_{\mu\nu} R^{(g)} - R_{\mu\nu}^{(g)} \right)$ and Eq. (92), we obtain

$$0 = -g_{\mu\nu} \nabla_g^\mu \left(\text{tr} \sqrt{g^{-1}f} \right) + \frac{1}{2} \nabla_g^\mu \left\{ f_{\mu\rho} \left(\sqrt{g^{-1}f} \right)^{-1\rho}{}_\nu + f_{\nu\rho} \left(\sqrt{g^{-1}f} \right)^{-1\rho}{}_\mu \right\}. \quad (93)$$

Similarly by using the covariant derivative ∇_f^μ with respect to the metric f , from (91), we obtain

$$0 = \nabla_f^\mu \left[\sqrt{\det(f^{-1}g)} \left\{ -\frac{1}{2} \left(\sqrt{g^{-1}f} \right)^{-1\nu}{}_\sigma g^{\sigma\mu} - \frac{1}{2} \left(\sqrt{g^{-1}f} \right)^{-1\mu}{}_\sigma g^{\sigma\nu} + \det \left(\sqrt{g^{-1}f} \right) f^{\mu\nu} \right\} \right]. \quad (94)$$

In case of the Einstein gravity, the conservation law of the energy-momentum tensor depends from the Einstein equation. It can be derived from the Bianchi identity. In case of bigravity, however, the conservation laws of the energy-momentum tensor of the scalar fields are derived from the scalar field equations. These conservation laws are independent of the Einstein equation. The Bianchi identities give equations (93) and (94) independent of the Einstein equation.

We now assume the FRW universes for the metrics $g_{\mu\nu}$ and $f_{\mu\nu}$ and use the conformal time t for the universe with metric $g_{\mu\nu}$:

$$ds_g^2 = \sum_{\mu,\nu=0}^3 g_{\mu\nu} dx^\mu dx^\nu = a(t)^2 \left(-dt^2 + \sum_{i=1}^3 (dx^i)^2 \right),$$

$$ds_f^2 = \sum_{\mu,\nu=0}^3 f_{\mu\nu} dx^\mu dx^\nu = -c(t)^2 dt^2 + b(t)^2 \sum_{i=1}^3 (dx^i)^2. \quad (95)$$

F(R) bigravity: Cosmological Reconstruction and Cosmic Acceleration

Then (t, t) component of (90) gives

$$0 = -3M_g^2 H^2 - 3m^2 M_{\text{eff}}^2 (a^2 - ab) + \left(\frac{3}{4} \dot{\varphi}^2 + \frac{1}{2} V(\varphi) a(t)^2 \right) M_g^2, \quad (96)$$

and (i, j) components give

$$0 = M_g^2 (2\dot{H} + H^2) + m^2 M_{\text{eff}}^2 (3a^2 - 2ab - ac) + \left(\frac{3}{4} \dot{\varphi}^2 - \frac{1}{2} V(\varphi) a(t)^2 \right) M_g^2. \quad (97)$$

Here $H = \dot{a}/a$. On the other hand, (t, t) component of (91) gives

$$0 = -3M_f^2 K^2 + m^2 M_{\text{eff}}^2 c^2 \left(1 - \frac{a^3}{b^3} \right) + \left(\frac{3}{4} \dot{\xi}^2 - \frac{1}{2} U(\xi) c(t)^2 \right) M_f^2, \quad (98)$$

and (i, j) components give

$$0 = M_f^2 (2\dot{K} + 3K^2 - 2LK) + m^2 M_{\text{eff}}^2 \left(\frac{a^3 c}{b^2} - c^2 \right) + \left(\frac{3}{4} \dot{\xi}^2 - \frac{1}{2} U(\xi) c(t)^2 \right) M_f^2. \quad (99)$$

Here $K = \dot{b}/b$ and $L = \dot{c}/c$. Both of Eq. (93) and Eq. (94) give the identical equation:

$$cH = bK \text{ or } \frac{c\dot{a}}{a} = \dot{b}. \quad (100)$$

If $\dot{a} \neq 0$, we obtain $c = \dot{a}b/\dot{a}$. On the other hand, if $\dot{a} = 0$, we find $\dot{b} = 0$, that is, a and b are constant and c can be arbitrary.

F(R) bigravity: Cosmological Reconstruction and Cosmic Acceleration

We now redefine scalars as $\varphi = \varphi(\eta)$ and $\xi = \xi(\zeta)$ and identify η and ζ with the conformal time t , $\eta = \zeta = t$. Hence, one gets

$$\omega(t)M_g^2 = -4M_g^2(\dot{H} - H^2) - 2m^2M_{\text{eff}}^2(ab - ac), \quad (101)$$

$$\tilde{V}(t)a(t)^2M_g^2 = M_g^2(2\dot{H} + 4H^2) + m^2M_{\text{eff}}^2(6a^2 - 5ab - ac), \quad (102)$$

$$\sigma(t)M_f^2 = -4M_f^2(\dot{K} - LK) - 2m^2M_{\text{eff}}^2\left(-\frac{c}{b} + 1\right)\frac{a^3c}{b^2}, \quad (103)$$

$$\tilde{U}(t)c(t)^2M_f^2 = M_f^2(2\dot{K} + 6K^2 - 2LK) + m^2M_{\text{eff}}^2\left(\frac{a^3c}{b^2} - 2c^2 + \frac{a^3c^2}{b^3}\right). \quad (104)$$

Here

$$\omega(\eta) = 3\varphi'(\eta)^2, \quad \tilde{V}(\eta) = V(\varphi(\eta)), \quad \sigma(\zeta) = 3\xi'(\zeta)^2, \quad \tilde{U}(\zeta) = U(\xi(\zeta)). \quad (105)$$

Therefore for arbitrary $a(t)$, $b(t)$, and $c(t)$ if we choose $\omega(t)$, $\tilde{V}(t)$, $\sigma(t)$, and $\tilde{U}(t)$ to satisfy Eqs. (101-104), the cosmological model with given $a(t)$, $b(t)$ and $c(t)$ evolution can be reconstructed. Following this technique we presented number of inflationary and/or dark energy models as well as unified inflation-dark energy cosmologies. The method is general and maybe applied to more exotic and more complicated cosmological solutions.

What is the next?

What is the next? So far $F(R)$ gravity which also admits extensions as HL or massive gravity is considered to be the best: simplest formulation, ghost-free, easy emergence of unified description for the universe evolution, friendly passing of cosmological bounds and local tests, absence of singularities in some versions (Bamba-Nojiri-Odintsov 2007), possibility of easy further modifications. More deep cosmological tests are necessary to understand if this is final phenomenological theory of universe and how it is related with yet to be constructed QG!